



NATUREX is a natural biofungicide based on Zinc, Tea tree (Melaleuca alternifolia ) oil and terpenic alcohols which prevents oxidative stress, and in particular, the damage caused by different fungi.

**NATUREX** contains Melaleuca Alternifolia Tea Tree Oil and anti-disease agents as well as other adjuvants to ensure nutrient uptake and improve product efficacy.

NATUREX is a natural Biofungicide that acts in a preventive and curative way, by inhibiting the development of spore germination, inhibition of mycelial growth and expansive lesion; inhibition in the production of sporangia, by suppression and eradication of colonies of pathogens present in fruits and leaves.

With its unique mode of action, NATUREX is an excellent tool for Resistance Management. It can be applied throughout the year without resistance and is non-toxic to crops, users and the environment.



Black sigatoka

**Powdery mildew** (Mycosphaerella fijiensis) (Erysiphe cichoracearum) (Botrytis cinerea



- Multiple modes of action
- Control of a wide range of plant pathogens, particulary bacterial and ascomycete diseases

## **COMPOSITION** %w/w

Melaleuca Alternifolia extract	20,0
Vegetable Oils	76,0
Zinc (Zn)	1,5
Manganese (Mn)	0,5



- No measurable affect on beneficial insects and bees
- Easily adapted sustaintable and IPM practices

## **DOSES AND APPLICATION**

Crop	Disease	Latin name	Dose ml/Ha	Crop	Disease	Latin name	Dose ml/Ha
Bananas	Black sigatoka	Mycosphaerella fijiensis	350-900	Rice	Blast	Pyricularia oryzae	500-1000
	Yellow sigatoka	Mycosphaerella musicola			Grain complex	Bipolaris oryzae	
Berries	Alternaria		800-1500		Sheath blight		
	Anthracnose			Tomato	Bacterial diseases		800-1500
	Fruit rot					Xanthomonas spp.	
	Grey mold						
	Powdery mildew						
Cucurbits	Powdery mildew		800-1500		Early blight		
					Grey mold		
Leafy greens	Powdery mildew		500-1000		Leafmold		
	White mold				Powdery mildew		
Peanuts	Early leaf spot		800-1000				
Peppers	Alternaria rot		800-1500	Vines	Grey mold		800-1200
	Anthracnose				Powdery mildew		
	Bacterial canker				Sour rot complex		
	Bacterial spot				1 - C - C - C - C - C - C - C - C - C -		
	Frog-eye leaf spot			Wheat	Fusarium head		
	Grey mold				blight		
	Powdery mildew						
	Syringae leaf spot						

Apply NATUREX using suitable equipment to ensure thorough coverage of all foliage. A minimun of 200 L/Ha and a maximun of 1000L/Ha is recommended.

## **PACKING:**





